

KARAPETYAN, S.Ye.

Geometric values of some congruence invariants. Nauch. dokl.
vys. skoly; fiz.-mat. nauki no.1:48-52 '58. (MIRA 12:3)

1. Yerevanskiy gosudarstvennyy universitet im. Kh. Abovyana.
(Congruences (Geometry))

16(1)

AUTHOR: Karapetyan, S. V. E.

SOV/155-58-2-11/47

TITLE: Two Congruences With Common Invariants F and F' (Dve kongruentsii s obshchimi invariantami F i F')

PERIODICAL: Nauchnyye doklady vysshey shkoly. Fiziko-matematicheskiye nauki, 1958, Nr 2, pp 55-59 (USSR)

ABSTRACT: The present paper is a continuation of a recently published paper of the author [Ref 1]. There he considered transformations of tetrahedra and determined two invariants: F - invariant of the first focal surface (A_1) and F' - invariant of the second focal surface (A_2). In the present paper the author considers two congruences ($A_1 A_2$) and ($\bar{A}_1 \bar{A}_2$) between the rays of which there exists a one-to-one relation, for which the focal invariants of the corresponding focal surfaces of these congruences remain preserved.

There are 9 Soviet references.

ASSOCIATION: Yerevanskiy armyanskiy pedagogicheskiy institut (Yerevan Armenian Pedagogical Institute)

SUBMITTED: February 15, 1958

Card 1/1

AUTHOR: Karapetyan, S.Ye. SOV/20-122-3-3/57

TITLE: Harmonic Quadrics and Certain Congruence Ruled Surfaces (Garmonicheskkiye kvadriki i nekotoryye lineychatyye poverkhnosti kongruentsiy)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol 122, Nr 3, pp 335-338 (USSR)

ABSTRACT: The author considers congruences the harmonic ruled surfaces of which form quadrics. He applies Cartan's method according to the well-known work of Finikov [Ref 1,2] and essentially uses his own earlier results [Ref 3-5]. From these he takes the equations of the harmonic ruled surfaces of the first and second focal congruence surface and from these, in accordance with the requirement that the two harmonic ruled surfaces be quadrics he obtains certain invariant equations for characterizing the considered congruence. The properties of the considered congruences are now connected with the generated Laplace sequence - the configuration L.

Theorem: The two harmonic ruled surfaces of a focal surface are quadrics, if and only if this congruence generates the configuration L.

Theorem: Two harmonic quadrics of a congruence of L simul-

Card 1/3

Harmonic Quadrics and Certain Congruence Ruled
Surfaces

SOV/20-122-3-3/57

taneously are harmonic quadrics of the other congruences of this configuration.

Theorem: If the two harmonic ruled surfaces of the given focal surface of the congruence are quadrics, then all asymptotic ruled surfaces of this congruence are quadrics too.

Theorem: If the two asymptotic ruled surfaces of a focal surface of the congruence are quadrics and if the configuration L generated by the congruence closes after the fourth step, then the harmonic ruled surfaces of this congruence are quadrics too.

In the last section of the paper some further remarkable ruled surfaces of the congruence are given and their properties are considered.

There are 6 Soviet references.

ASSOCIATION: Armyanskiy gosudarstvennyy pedagogicheskiy institut imeni Kh.Abovyana (Armenian State Pedagogical Institute imeni Kh. Abovyan)

PRESENTED: April 25, 1958, by S.L. Sobolev, Academician

Card 2/3

KARAFETI, A.S. 1/2

16(1)
 PHASE I BOOK EXPLOITATION
 SOV/2660
 Vsesoyuzny matematicheskiy s'yezd. 3rd, Moscow, 1956
 Trudy. t. 4: Krestnye soderzhanie sektsionnykh dokladov. Doklady
 izrazheniya uchastnikov (Transactions of the 3rd All-Union Mathema-
 tical Conference, 1. Moscow, vol. 4: Summary of Sectional Reports.
 Reports of Foreign Scientists) Moscow, Izd-vo AN SSSR, 1959.
 247 p. 2,200 copies printed.
 Sponsoring Agency: Akademiya nauk SSSR. Matematicheskiy Institut.
 Tech. Ed.: G.M. Shevchenko; Editorial Board: A.A. Abramov, V.G.
 Boltzanskiy, A.M. Vasil'yev, B.V. Medvedev, A. Rybnik, S.M.
 Mikul'skiy (resp. Ed.), A.G. Postnikov, Yu. G. Izrael, E.M.
 Rybnikov, P. L. Vilyanov, V.A. Uspenskiy, N.G. Chetayev, O. Ya.
 Shilov, and A.I. Shiranov.

PURPOSE: This book is intended for mathematicians and physicists.
 COVERAGE: The book is Volume IV of the Transactions of the Third All-
 Union Mathematical Conference, held in June and July 1956. The
 book is divided into two main parts. The first part contains sum-
 maries of the papers presented by Soviet scientists at the Con-
 ference that were not included in the first two volumes. The
 second part contains the text of reports submitted to the editor
 by non-Soviet scientists. In those cases when the non-Soviet sci-
 entist did not submit a copy of his paper to the editor, the title
 of the paper is cited, and, to the extent possible, the title
 of the paper is included in the appropriate volume. The papers,
 both Soviet and non-Soviet, cover various topics in number theory,
 algebra, differential and integral equations, function theory,
 functional analysis, probability theory, topology, mathematical
 problems of mechanics and physics, computational mathematics,
 mathematical logic and the foundations of mathematics, and the
 history of mathematics.

Rybnikov, S.G. (Moscow). The invariance of infinite dimen-
 sional homology groups 73
 Section on Geometry
 Buzols, O.L. (L'viv). On certain problems of geometrography
 connected with accuracy of graphic computations 75
 Gerdaryk, D.Z. (Minsk). Incidence axioms of multidimen-
 sional projective geometry 75
 Kuz'man, A.G. (Stalingrad). Certain problems of local de-
 formability of surfaces 76
 Sazonov, S.Ye. (Yerevan). Linear complexes of developing
 surfaces of a congruence 76
 Lopatits, A.M. (Moscow). Fundamentals theorems of the theory
 of a hypersurface in dimensionless Euclidean space 77
 Card 15/ 34

3

16(1)

05689

AUTHOR:

Karapetyan, S.Ye.

SOV/22-12-4-2/9

TITLE:

The Pair A and Some Properties of the Pair T

PERIODICAL:

Izvestiya Akademii nauk Armyanskoy SSR. Seriya fiziko-matematicheskikh nauk, 1959, Vol 12, Nr 4, pp 27 - 34 (USSR)

ABSTRACT:

A configuration A of two congruences is defined by making correspond ruled surfaces to each ray of the congruences and then demanding that the tangential plane of each of these ruled surfaces passes through the point of tangency of the other ruled surface. The defined configuration forms a generalization of the T-configuration of Finikov [Ref 3,4]. The author proves some properties of the two configurations, e.g. the directions of the main ruled surfaces of T are identical with the main directions of the same configuration (see Calapso [Ref 6,7]). There are 12 Soviet references.

ASSOCIATION: Armyanskiy pedagogicheskiy institut imeni Kh.Abovyana
(Armenian Pedagogical Institute imeni Kh. Abovyan)

SUBMITTED:

October 27, 1958

Card 1/1

85237

S/022/60/013/002/008/011 XX
C 111/C222

14.5600

AUTHOR: Karapetyan, S.Ye.

TITLE: Configuration L

PERIODICAL: Izvestiya Akademii nauk Armyanskoy SSR. Seriya fiziko-matematicheskikh nauk, 1960, Vol. 13, No. 2, pp. 3 - 16.

TEXT: The present paper joins the author's earlier papers (Ref. 3,4,10,11) and uses the same notations in (Ref. 3) the author obtained the equation of the Lie quadric for all ruled surfaces

$\left(\omega_2^4 = \lambda \omega_1^3 \right)$ of the congruence. On the first focal surface ω to their harmonic ruled surfaces there corresponds the conjugate net $(\alpha(\omega_1^3)^2 - \gamma(\omega_2^4)^2 = 0)$ with the property that both tangents of the net form a harmonic quadruple with the two tangents of the first focal net. Thus the mentioned conjugate net is called the harmonic net Γ_1 of the first focal surface (A_1) . The harmonic ruled surfaces of the congruence

Card 1/4

85237

Configuration L

S/022/60/013/002/008/011 XX
C111/C222

(A_1, A_2) which correspond to the net Γ_1 are again denoted with $\Gamma_1 \cdot \Gamma_2$ is the harmonic net of the focal surface (A_2) etc.

Amongst other things it is shown : Two focal surfaces of the congruence and two tangential planes of two arbitrary ruled surfaces of the congruence (with a common tangential point) form a harmonic quadruple then and only then if these ruled surfaces are conjugate in the sense of Sania. (A_1) is

the common focal surface of two congruences of the Laplace sequence. Ruled surfaces belonging to an other congruence of the sequence and corresponding to the lines Γ_1 are denoted with Γ'_1 . The Lie quadrics of $\Gamma_1, \Gamma_2, \Gamma'_1, \Gamma'_2$ are determined. Two geometric characteristics of two ruled surfaces,

conjugate in the sense of Sania, of the congruence are given. If a pair of corresponding quadrics Γ_1 and Γ'_1 has a common tangential plane

along the ray of the given congruence, then it has a common tangential plane along the ray of the following congruence of the Laplace sequence. If in each pair the corresponding quadrics of Γ_1 and Γ'_1 coincide, then

Card 2/ 4

85237

Configuration L

S/022/60/013/002/008/011 XX
C111/C222

it holds : 1) The sequence generated by the congruence (A_1, A_2) is an R-sequence (cf. (Ref. 6)); 2) Each pair of corresponding quadrics of Γ_2 and Γ'_2 coincides; 3) All focal surfaces of the sequence are surfaces of second order; 4) Such a configuration is denoted as a configuration L and depends on 10 arbitrary constants. This configuration L is formed by two surfaces of second order. The diagonals A_1A_4 and A_2A_3 are simultaneously conjugate with respect to both surfaces and describe a congruence of straight lines each directrix of which intersects all tangents of a number of the nets Γ_1 and Γ_2 .

There are 11 references : 10 Soviet and 1 Italian.

[Abstracter's note : (Ref. 3,4,10,11) are papers of the author in Doklady Akademii nauk SSSR 1957, Vol. 117, No. 2; Nauchnyye doklady vysshey shkoly. Fiziko-matematicheskkiye nauki, 1958, No. 1; Nauchnyye doklady vysshey shkoly. Fiziko-matematicheskkiye nauki, 1958, No. 2; Doklady

Card 3/4

85237

84303

16.5600

S/022/60/013/004/005/007XX
C111/C222

AUTHOR: Karapetyan, S.Ye.

TITLE: On a Congruence Transformation 16

PERIODICAL: Izvestiya Akademii nauk Armyanskoy SSR. Seriya fiziko-matematicheskikh nauk, 1960, Vol.13, No.4, pp.3-17

TEXT: The papers (Ref.1,2,3,4) assert that the straight lines of the P_3 are mapped onto the points of the hyperquadratic Q_4^2 of the P_5 . The tangential plane intersects Q_4^2 in a three-dimensional cone K_3^2 the vortex of which lies in the tangential point. The asymptotic lines on Q_4^2 are determined by the differential equation

$$(5) \quad \omega_1^3 \omega_2^4 - \omega_2^3 \omega_1^4 = 0.$$

The quadratic fundamental form (5) defines a polarity of the varieties in the tangential hyperplane. Two varieties L_m and L_{m-4} being incident to the tangential point and to the tangential plane, are called conjugated if they correspond to each other according to this polarity. In the preset paper the author uses the same conjugate varieties in order

Card 1/4

S/022/60/013/004/005/007XX
C111/C222

On a Congruence Transformation

to obtain a congruence transformation and to obtain some theorems. The author uses the method of Cartan according to (Ref.5). The conjugate varieties can be described as follows: Let the two-dimensional surface (p_1) be the image of the congruence in the P_5 and L_1 be the tangent of (p_1) along the line $\omega_2^4 = \lambda \omega_1^3$. In the polarity (5), to the straight line L_1 there corresponds a certain L_3 which intersects the tangential 2-plane of (p_1) in a certain L_1' . The directions L_1 and L_1' are called conjugate in the congruence (P_1) . The L_3 -characteristic along L_1 is the 2-plane L_2 having only one common point p with the tangential 2-plane of (p_1) . An analogous point p' can be obtained if L_1 and L_1' are changed. The construction of the point p and p' in the P_3 leads to Laplace transformations. The straight line pp' intersects Q_4^2 in certain given points p_1' and p_1'' . In the P_3 every straight line P_1' , P_1'' lies on a focal surface and goes through the other focus of the congruence (P_1) . The (p_1') and (p_1'') are denoted as Π -transformations of the congruence

Card 2/4

84303

S/022/60/013/004/005/007XX
C111/C222

On a Congruence Transformation

(p_1) in the conjugated directions $\omega_2^4 = \pm \lambda \omega_1^3$. The author gives conditions for the fact that the transformations Π and the focal surfaces of (p_1) are harmonic and conjugated, respectively. The case where the developable surfaces of the congruences (p_1') and (p_1'') intersect with the focal surfaces (p_1) along asymptotic lines, is investigated separately. Furthermore the case where the developable surfaces of the congruences (p_1') and (p_1'') intersect with the focal planes of the congruence (P_1) in a harmonic net is investigated. In this case (p_1) , (p_1') , (p_1'') are W-congruences and the asymptotic lines on their focal surfaces correspond to each other. Finally the case is investigated, where the straight lines p_1' and p_1'' are polarly conjugated with respect to the two Lie quadrics of the focal surfaces of

Card 3/4

KARAPETYAN, S.Ye.

Quadrics of congruences. Dokl. AN Arm. SSR 30 no.2:
65-72 '60. (MIRA 13:6)

1. Yerevanskiy armyanskiy pedagogicheskiy institut imeni
Kh. Abovyana. Predstavleno akad. AN Armyanskoy SSR
M.M. Dzhrbashyanom.
(Congruences(Geometry))

KARAPETYAN, S.Ye. (Yerevan)

Transformation of congruences by means of semiquadrics. Mat.shor.
50 no.1:109-116 Ja '60. (MIRA 13:6)
(Congruences)

06/13/2000

CIA-RDP86-00513R000720610017-7"

S/020/60/133/005/022/034XX
C111/C222

16.5600

AUTHOR: Karapetyan, S.Ye. 16

TITLE: Conjugate Manifolds

PERIODICAL: Doklady Akademii nauk SSSR, 1960, Vol.133, No.5, pp.1007-1010.
TEXT: As it is well-known (Ref.2-4), the straight lines of the P_3 are mapped into points of the hyperquadric Q_4^2 of the P_5 . Q_4^2 generates a polarity of the linear subspaces in the P_5 , where the sum of dimensions of the conjugate spaces is 4. If a subspace lies in the tangential hyperplane of the Q_4^2 and passes through the tangential point, then the conjugate subspace has this property too. Such subspaces are called conjugate manifolds. The common subspace of two conjugate manifolds L_m and L_{4-m} is called an asymptotic manifold.
An infinitesimal displacement of the tetrahedron $A_1 A_2 A_3 A_4$ in the projective space is described by
(1) $da_i = \omega_i^k A_k$, $i, j, k=1, 2, 3, 4$.

84643

S/020/60/133/005/022/034XX
C111/C222

Conjugate Manifolds and Their Application

where ω_i^k are linear differential forms connected with the structural equations of the space $D\omega_i^k = [\omega_i^j \omega_j^k]$ (cf. (Ref.1)). Let the analytic straight lines be denoted by

$$(2) \quad p_1 = (A_1 A_2), \quad p_2 = (A_3 A_4), \quad p_3 = (A_2 A_3), \quad p_4 = (A_1 A_4), \quad p_5 = (A_1 A_3),$$

$$p_6 = (A_4 A_2).$$

The tangential hyperplane of the Q_4^2 in the point p_1 is determined by the Grassmann product $(p_1 p_3 p_4 p_5 p_6)$. The asymptotic manifold is determined from the postulate $(d^2 p_1 p_3 p_4 p_5 p_6 p_1) = 0$, which leads to

$$(3) \quad \omega_1^3 \omega_2^4 - \omega_2^3 \omega_1^4 = 0.$$

If $A_1 A_2$ describes a ruled surface L in P_3 , then its image describes a curve l on Q_4^2 ; if the ruled surface is developable, then the image

Card 2/3

KARAPETYAN, S.Ye.

Transformation of a congruence by means of ruled surfaces.

Dokl. AN Arm. SSR 32 no.1:9-16 '61.

(MIRA 14:3)

1. Armyanskiy pedagogicheskiy institut imena Kh.Abovyana.
Predstavleno akademikom AN Armyanskoy SSR M.M. Dzhrbashyanom.
(Congruences(Geometry)) (Surfaces, Ruled)

KARAPETYAN, S.Ye.

Theory of pairs of congruences. Izv. AN Arm. SSR. Ser. fiz.-mat.
nauk 14 no.4:37-47 '61. (MIRA 14:11)

1. Armyanskiy pedagogicheskiy institut imeni Kh. Abovyana.
(Congruences (Geometry))

KARAPETYAN, S.Ye.

Linear manifolds of straight lines and planes in four-dimensional projective space. Izv.AN ~~Arm.~~ SSR.Ser.fiz.-mat.nauk 15 no.1:53-72 '62. (MIRA 15:2)

1. Armyanskiy pedagogicheskiy institut imeni Kh.Abovyana i institut matematiki i mekhaniki AN Armyanskoy SSR.
(Geometry, Projective)

KARAPETYAN, S.Ye. (Yerevan)

Complexes of straight lines in three-dimensional projective space. Mat.sbor. 56 no.3:343-352 Mr '62. (MIRA 15:4)

1. Armyanskiy pedagogicheskiy institut imeni Kh.Abovyana.
(Aggregates) (Geometry, Differential)

KARAPETYAN, S.Ye.

Projective-differential geometry of a two-parameter family of straight lines and planes in a four-dimensional space. Part 1.
Izv.AN Arm.SSR.Ser.fiz.-mat.nauk 15 no.2:25-43 '62. (MIRA 15:4)

1. Armyanskiy pedagogicheskiy institut imeni Kh. Abovyana i
Institut matematiki i mekhaniki AN Armyanskoy SSR.
(Geometry, Differential--Projective)

KARAPETYAN, S.Ye.

Projective differential geometry of two-parameter families
of straight lines and surfaces in four-dimensional space.
Izv. AN Arm. SSR. Ser. fiz.-mat. nauk 15 no.3:17-28 '62.

(MIRA 15:9)

1. Armyanskiy pedagogicheskiy institut imeni Kh. Abovyana
i Institut matematiki i mekhaniki AN Armyanskoy SSR.
(Geometry, Differential--Projective)

KARAPETYAN, S.Ye.

Projective differential geometry of families of n -dimensional planes. Part 1. Izv. AN Arm SSR. Ser. fiz.-mat. nauk 16 no.3: 3-22 '63. (MIRA 16:8)

1. Yerevanskiy zaachnyy pedagogicheskiy institut i Institut matematiki i mekhaniki AN Arm SSR.
(Geometry, Differential—Projective)

KARAPETYAN, S.Ye.

Projective differential geometry of a family of multidimensional planes. Part.2. Izv. AN Arm.SSR.Ser.fiz.-mat. nauk 16 no.5:3-22 '63. (MIRA 16:11)

1. Institut matematiki i mekhaniki AN Armyanskoy SSR i Armyanskiy zaochnyy pedagogicheskiy institut.

KARAPETYAN, S.Ye.

Projective differential geometry of families of n -dimensional planes.
Part 3. Izv. AN Arm. SSR.Ser.fiz.-mat.nauk 17 no.1:3-21 '64.

(MIRA 17:3)

1. Armyanskiy zaachnyy pedagogicheskiy institut i Institut matematiki
i mekhaniki AN Armyanskoy SSR.

KARAPETYAN, T.A.

Distribution and prospects for the development of the industry of mining building materials in the Shiraki Steppe [in Armenian with summary in Russian]. Nauch. trudy Yrev. un. 63:183-208. '58.

(MIRA 11:6)

1. Yerevanskiy gosudarstvennyy universitet, kafedra ekonomicheskoy geografii.

(Armenia—Building materials industry)

AID P - 4937

Subject : USSR/Electronics

Card 1/1 Pub. 89 - 4/18

Author : Karapetyan, V.

Title : Ultrashort wave school radio stations

Periodical : Radio, 8, 19-20, Ag 1956

Abstract : The author points to the necessity of paying more attention to the development of ultrashort wave radio stations in schools. These school stations contribute to the familiarization of more people with radio and radio equipment.

Institution : None

Submitted : No date

GUKASYAN, V.; KARAPETYAN, V.

Production of concrete with a natural texture finish for outer wall panels. Prom. Arm. 5 no. 8:54-55 Ag '62. (MIRA 15:8)

1. Armyanskiy institut stroitel'nykh materialov i sooruzheniy.
(Armenia—Building materials)

KARAPETYAN, V.A.

Brief reports. Zav. lab. 25 no.1:126 '59.

(MIRA 12:1)

1. Moskovskiy aviatsionny tekhnologicheskoy institut.
(Grinding machines)

KARAPETIAN, V.A.

33141

S/120/61/000/006/007/041
E032/E114

21.6000

AUTHORS: Khrimyan, A.V., Yeghyan, K.Sh., Nalbandyan, N.A.,
Avakyan, V.V., and Karapetian V.A.

TITLE: Measurement of charged-particle masses with the aid
of scintillation counters

PERIODICAL: Priory i tekhnika eksperimenta, no.6, 1961, 52-56

TEXT: The method can be used to (a) select particles which
stop in the scintillator owing to ionization losses, and
(b) to determine the mass of the particles by measuring their
energy and range in the scintillator. The device consists of a
telescope of n scintillation counters (C_1, \dots, C_n) with
thickness l_1, \dots, l_n respectively. If a particle which has
passed at an angle of φ through $k - 1$ scintillators has come
to rest in the scintillator C_k at a depth l_x , and at the end
of its range in the $m + 1$ scintillators C_{k-m}, \dots, C_k the
energy losses $\Delta E_{k-m}, \dots, \Delta E_k$ were due to ionization only, then
it can be shown that:

Card 1/6

X

Measurement of charged-particle ...

33311
S/120/61/000/006/007/041
EO32/E114

$$\frac{\Delta E_{k-i}}{\Delta E_{k-(i+1)}} = f_i \left(\frac{\Delta E_{k-(i+1)}}{\Delta E_{k-(i+2)}}, \ell_{k-1}, \dots, \ell_{k-(i+2)} \right) \quad (1)$$

(i = 0, ..., m - 2)

This holds whatever the nature of the particle, the direction of its motion, and range in the last scintillator C_k . Thus, by measuring the energies $\Delta \mathcal{E}_1, \dots, \Delta \mathcal{E}_n$ in the scintillators C_1, \dots, C_n one can select with the aid of Eq.(1) all those particles which come to rest in the scintillators C_{k-m}, \dots, C_k by losing energy in ionization processes only. For stable particles $\Delta \mathcal{E}_i = \Delta E_i$. If on the other hand a primary particle decays (or is captured) in the scintillator C_k then the energy liberated in C_k is $\Delta \mathcal{E}_k = \Delta E_k + \delta E_k$ where the latter quantity is the energy of the secondary particles. In this case the first equation (i = 0) in Eq.(1) can only be used for the determination of the unknown energy:

$$\Delta E_k = \Delta \mathcal{E}_{k-1} f_0 (\Delta \mathcal{E}_{k-1} / \Delta \mathcal{E}_{k-2}) \quad (4)$$

Card 2/6

331h1

Measurement of charged-particle...

S/120/61/000/006/007/041
E032/E114

and the remaining relations in Eq.(1) are used to select the ionization stoppages. The energy loss of a particle with an ionizing power $1/I_{\min}$ in the scintillator C_1 is given by:

$$\Delta \xi_i = B (I/I_{\min})_{C_1} \ell_i \text{ MeV} \quad (5)$$

where B is in MeV/cm and represents the minimum ionization loss in the particular scintillator, and ℓ_i is the thickness of the scintillator C_i in cm. Thus the energy lost by a particle before stopping in scintillators C_{k-m}, \dots, C_k is given by

$$E = \sum_{i=k}^{k-m} \Delta \xi_i$$

If Eq.(1) is not satisfied for $i = 0$, then

$$E = \sum_{i=k-1}^{k-m} \Delta \xi_i + \Delta E_k \quad (6)$$

Card 3/6

Measurement of charged-particle...

333h1
S/120/61/000/006/007/041
E032/E114

where ΔE_k is given by Eq.(4). The range of a particle in the scintillators C_{k-m}, \dots, C_k is given by:

$$R = \left(\sum_{i=k-1}^{k-m} \ell_i + \ell_x \right) \operatorname{cosec} \varphi \quad (7)$$

in which all the quantities except ℓ_x are known. If the scintillators are looked upon as simple filters then

$$\ell_x = 1/2 \ell_k \pm 1/2 \ell_k.$$

ℓ_x can also be determined from a relation of the form:

$$\ell_x = F(\ell_0, \ell_{k-1}, \ell_{k-2}) \quad (3)$$

In order to verify the above method the authors have used the results obtained with the instrument described by A.I. Alikhanov, A.V. Khrimyan, V.K. Kosmachovskiy, V.V. Avakyan, Yu.V. Gorodkov, K.Sh. Yeghyan and N.A. Nalbandyan (Ref.6: Proceedings of the International Conference on Cosmic Rays, 1959, 1960, v.1, 183)

Card 4/6

Measurement of charged-particle ...

33141
S/120/61/000/006/007/041
E032/E114

The instrument consists of a magnetic mass spectrometer, a five-layer proportional counter (A.I. Alikhanov, V.A. Lubimov, G.P. Elisiyev, CERN Symposium, v.2, 1956, 87) and five scintillation counters (V.K. Kosmachevskiy and M.S. Aynuddinov, PTE, no.3, 1956, 49). The rms error in the momenta between 0.2 and 1 GeV/c was approximately 8 to 5% for protons and 2 to 4% for π -mesons. The ionizing power of the particles could be measured with the proportional counter to an average accuracy of $\pm 14\%$. For particles stopping in the scintillation counters the average losses in the scintillators could be measured to $\pm 10\%$. Preliminary results indicate that the efficiency of selection of particles which come to rest owing to ionization only is about 0.8. The average accuracy with which the masses can be determined from the energies and ranges is approximately 20%. The statistics on which these results are based are limited and therefore the results are only preliminary. The experiment did not confirm the possibility of investigating the masses and decays of unstable particles. The method may find wide-ranging applications and is amenable to automation. Acknowledgments are

Card 5/6

Measurement of charged-particle ...

3341
S/120/61/000/006/007/041
E032/E114

expressed to A.I. Alikhanov and A.I. Alikhanyan for interest and discussions, and to Yu.V. Gorodkov, M.P. Lorikyan, I.P. Karabekov, K.A. Khurshudyan, G.P. Matevosyan, V.V. Truzyan, E.V. Patvakanyan, G.M. Smsaryan, A.A. Oganessian and B.V. Tovmasyan for assistance in the organisation and execution of this work.

There are 4 figures and 11 references; 5 Soviet-bloc and 6 non-Soviet-bloc. The four most recent English language references read as follows:

Ref.2: J.W. Keuffel, R.L. Call, W.H. Sandmann, M.O. Larson.
Phys. Rev. Letters, v.1, 1958, 203.

Ref.4: Phys. Rev., v.114, 1959, 1150.

Ref.5: E. Birman, R. Lea, J. Orear, S. Rosendorff.
Phys. Rev., v.113, 1959, 710.

Ref.7: J. Steinberger, 1958 Annual International Conference on
High Energy Physics at CERN, Geneva, 1958.

ASSOCIATION: Fizicheskii institut AN ArmSSR
(Physics Institute, AS Armenian SSR)

SUBMITTED: April 3, 1961
Card 6/6

35969

S/048/62/026/006/019/020
B125/B102

9.6150

AUTHORS: Khrimyan, A. V., Yeghyan, K. Sh., Nalbandyan, N. A.,
Avakyan, V. V., and Karapetyan, V. A.

TITLE: On the measurement of masses of charged particles by means
of scintillation counters

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya fizicheskaya,
v. 26, no. 6, 1962, 831-836

TEXT: A group of scintillation counters can be used to determine the
stoppings due to ionization losses and the masses (range-energy
measurement). The apparatus here used comprised a magnetic mass spectro-
meter ($H = 6850$ oe), a five-layer proportional counter and five
scintillation counters. After measuring the energies released from the
particle in the scintillators C_1, \dots, C_n with the thicknesses
 l_1, \dots, l_n ($n \geq 3$) the stoppings due to ionization losses were
distinguished from the nuclear interactions by applying the criterion

Card 1/3

On the measurement of masses ...

S/048/62/026/006/019/020
B125/B102

$$\frac{\Delta E_{k-i}}{\Delta E_{k-(i+1)}} = f_i \left(\frac{\Delta E_{k-(i+1)}}{\Delta E_{k-(i+2)}}, l_{k-1}, \dots, l_{k-(i+2)} \right) \quad (i=0, \dots, m-2)$$

$\Delta E_{k-m}, \dots, \Delta E_k$ are the energy losses in the scintillators

C_{k-m}, \dots, C_k . The four quantities momentum, ionization power, range and energy are measured by this device. From these, the mass of the particles is found by the momentum - ionization and range - energy methods. The mass spectrum as measured by the first method has a maximum at $\sim 1780 m_e$ and that obtained from the second method a maximum at $\sim 1850 m_e$.

In both cases a weak deuteron spectrum appears between 3500-4500 m_e . The stoppings due to ionization are identified with an efficiency of ~ 0.8 . The stoppings due to other causes are eliminated with an efficiency of $\sim 0.9-1$. This method was tested by the devices available at the time and can undoubtedly be improved upon by more perfect selection and use of apparatus. Its applicability to decay processes and to mass measurements of unstable particles has not yet been confirmed experimentally. There are 4 figures. The most important English-language reference is: Stenberger J. 1958 Annual International Conference on High Energy Physics at CERN, Geneva, 1958.

Card 2/3

S/048/62/026/006/020/020
B181/B104

AUTHORS: Khrimyan, A. V., Yegiyany, K. Sh., Nalbandyan, N. A.,
Avakyan, V. V., and Karapetyan, V. A.

TITLE: Mass measurements of low-intensity charged-particle groups
by various methods

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya fizicheskaya, v. 26,
no. 6, 1962, 837- 840

TEXT: The mass of particles produced by the action of cosmic rays was determined from (1) momentum and ionization, (2) momentum and length of path, (3) momentum and energy, (4) ionization and energy, (5) ionization and length of path, (6) energy and length of path. The experimental arrangement (A. V. Khrimyan, V. V. Avakyan, N. A. Nalbandyan, K. Sh. Yegiyany, M. P. Pleshko, present publication, p. 722) consisted of a mass spectrometer, a proportional counter, two scintillation counters for determining the energy and length of path, and three scintillation counters for determining the energy losses of scattered particles. (2) and (3) gave masses too high, (4), (5); and (6) masses too small for the 203

Card 1/2

KHRIMYAN, A.V.; YEGIYAN, K.Sh.; NALBANDYAN, N.A.; AVAKYAN, V.V.; KARAPETYAN, V.A.

Measurement of the masses of charged particles with the aid of scintillation counters. Izv. AN SSSR. Ser. fiz. 26 no.6:831-836 Je '62. (MIRA 15:6)

1. Fizicheskiy institut Akademii nauk Armyanskoy SSR.
(Scintillation counters) (Particles (Nuclear physics))

KRIMYAN, A.V.; YEGIYAN, K.Sh.; NALBANDYAN, N.A.; AVAKYAN, V.V.; KAPAPETYAN,
V.A.

Various methods for measuring the masses of low-intensity groups of
charge particles. Izv. AN SSSR. Ser. fiz. 26 no.6:837-840 Je '62.
(MIRA 15:6)

1. Fizicheskiy institut Akademii nauk Armyanskoy SSR.
(Particles (Nuclear physics)) (Mass spectrometry)

AKOPYAN, A.V.; KARAPETYAN, V.A.

Experimental study of the rigidity of reinforced tiffcrete beams under the prolonged action of loads. Izv. AN Arm. SSR. Ser. tekhn. nauk 17 no. 4:77-82 '64. (MIRA 17:11)

1. Armyanskiy nauchno-issledovatel'skiy institut stroitel'nykh materialov i sooruzheniy.

KARAPETYAN, V. K.

PA 43/473

"Agrobiologiya" No 4

Changes in life characteristics of wheat result in a series of changes and morphological indices. Conducted experiments to determine changes of hard wheat into soft wheat. Observed most noticeable changes in vernalized hard wheat. Noticeable changes in season yielded soft wheat when sown during fall season's theory that wheat based experiments on Lyenko's theory if left in the ground in winter. Many tables show experimental results.

43/4913

USSR/Agriculture
Plant Breeding
Wheat

Jul/Aug 48

"Changing the Nature of Hard Wheats Into Soft
Wheats," V. K. Karapetyan, Inst of Genetics, Acad
Sci USSR, 17 pp

KARAPETTYAN, V. K.

PA 43/49T72

USSR/Medicine - Plants
Medicine - Environment

Jul/Aug 48

"Possibility of Transforming Certain Types of
Cultivated Plants Into Other Types" 2 pp

"Agrobiologiya" No 4

Neo-Darwinist-Morganist theory of nonhereditary
phenotypic changes has been disapproved. Michurin's
theory of genetics is the right one, and its
acceptance will result in greater achievements by
Soviet biologists and selection specialists. Re-
fers to work by V. K. Karapetyan, which proves
beyond all doubt that changes in the nature of an
organism are not hereditary factors but are
brought about by changes in its environment.

43/49T72

USSR/Medicine - Plants (Contd)

Jul/Aug 48

organism are not hereditary factors but are
brought about by changes in its environment.

43/49T72

KARAPETYAN, V. K.

Kar-petran, V. K. - "The softening of durum wheat," Yestestvo-zaniye v shkole, 1948, No. 6, 18--30

SO: U-3600, 10 July 53, (Letopis 'Zhurnal 'nykh Statey, No. 6, 1949).

110

Enzymic activity of hereditarily hard wheats changed to hereditarily soft wheats. N. M. Sisakyan, V. K. Karapet-
yan, and N. A. Vasil'eva (A.N. Bakh Binehem. Inst., Moscow). *Problemy Biokhim. i Molekul. Biol.*, Moscow, 1969, No. 1, 92-101 (1969); cf. C.A. 44, 5435g. — The dehydrogenase system of the altered wheat becomes close to that of naturally soft wheat; similarly the respiratory coeff. approaches the higher levels of the soft wheat. Polyphenoloxidase and peroxidase activities remain approx. const., while β -amylase approaches the levels found in soft wheat.
G. M. Kozolapoff

110

Direction of enzymic transformation of carbohydrates of hereditarily summer forms of wheat that had been altered to hereditarily winter forms. N. M. Sisakyan, V. K. Kataryan, and A. M. Kobayakova (A. N. Bakh Biokhem. Inst., Moscow), *Problemy Biokhim. i Molekul. Biol. Akad. Nauk S.S.S.R.*, Sbornik, No. 1, 102 (1969); cf. C.I. 44, 5435g. — Summer forms of wheat that were transformed by training into the winter forms display a shift in the enzymic transformations of sucrose; the enzymic action undergoes the adaptation of its synthetic action to the lower temp. and the high level (relative) of synthetic action approaches that found in natural winter varieties of wheat. G. M. K.

110

The inheritance of acquired enzymic characteristics of hard wheat which had been transformed into soft wheat. N. M. Sisakyan, V. K. Karapetyan, and N. A. Vasil'eva (Bach Biochem. (MIL. Moscow).—*Biokhimiya* 13, 9-13 (1930).—In biochem. properties, hard wheat differs from soft wheat in that the former possesses a greater dehydrogenase and β -amylase activity. Contrary to the teachings of Mendel and Morgan, hard wheat can be transformed into soft wheat by planting in an autumn-winter environment (*Agrobiologiya* 4, 5(1948)). Such soft wheat then also contains a lower dehydrogenase and β -amylase activity than the parent hard wheat. These acquired weaker enzymic characteristics are passed on to future generations.
H. Priestley

SAHN, V. K.

183T2

USSR/Biology - Genetics

Dec 50

"New Data on the Directed Alteration of Hard Wheat to Soft Wheat by Means of Winter Cultivation and Facts on the Transformation of Wheat into Rye," V. K. Karapetyan

"Trudy Inst Genetiki, Ak Nauk SSSR," No 18, pp 47-65

After repeated winter cultivation for several yr, grains of Tr. vulgare (I) develop on ears of Tr. durum (II). Plant then has 28 somatic chromosomes in some tissues (typical for II) and 42 in others (typical for I). In regions where conditions of wintering are very severe, individual grains of

LC

183T2

USSR/Biology - Genetics (Contd)

Dec 50

rye (Secale cereale) develop on plants of I and II. Such grains of rye, grown in complete isolation, develop into full-fledged rye plants which yield rye grains. Individual grains of wheat (1 per ear) were found on ears of wild fld rye of type which develops from rye grains originating on wheat. Such wheat grains had 42 chromosomes in dividing cells (typical for wheat), while all other grains of same ear had 14 chromosomes (typical for rye).

LC

183T2

KARAPEYATAN, V. K.

USSR/Biology, Agricultural Genetics

Mar/Apr 52

"Some New Data on the Transformation of Species in Cereal Plants," V. K. Karapetyan. Inst of Genetics, Acad Sci USSR

"Agrobiologiya" No 2, pp 29-44

Single wheat grains found in rye ears yielded 42-chromosome wheat plants; the remaining rye grains grew into typical 14-chromosome rye plants. Rye plants grown from rye grains found in ears of soft and hard wheat showed the same chromosome complement as ordinary rye plants. A new form of soft wheat which originated from hard wheat had 28 and

215T2

48 chromosomes in somatic tissues and 14 and 28 chromosomes in genetic cells, thus combining the properties of *Tr. durum* and *Tr. vulgare*. In some cases, grains resembling rye which originated from *Tr. durum* ears yielded spelt (*Tr. dicoccum*) plants with 28 chromosomes at the tips of the rootlets (a characteristic typical for spelt). Rye plants were obtained by planting vernalized ("yarovized") wheat grains either in the winter or spring (winter soft wheat and semi-winter hard wheat). The rye plants obtained were not hybrids: they were fully fertile. No intermediate forms between rye and wheat were observed. Expts proved that the rye grains found in wheat ears originated without fertilization. On castration and isolation,

215T2

plants from such grains also formed grains without fertilization. The rye plants from grains resulting on castration and isolation were quite normal and had 14 chromosomes at the tips of the rootlets, which is typical for rye.

215T2

ARAPETIAN, V. K.

2. USSR 600

4. Plants - Evolution

7. Species development in plants, Trudy Inst. gen, No. 19, 1952.

9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

ARAPETIAN, V. K.

Genetic analysis of the first and second generations of rye-wheat
and wheat-rye hybrids. Trudy Inst.gen. no.20:35-59 +53. (MLRA 7:1)
(Rye) (Wheat) (Hybridization, Vegetable)

Karapetyan, V.K.

U.S.S.R.

✓ Biochemical and technological characteristics of rye with wheat heredity. V. L. Kretovich, Ya. N. Kuprits, V. K. Karapetyan, and A. A. Bunder (A. N. Bakh Inst. Biochem. Sci. U.S.S.R., Moscow and Inst. Genetics, Acad. Sci. U.S.S.R.). *Biokhim. Zhurn., Akad. Nauk S.S.S.R., Sbornik 2*, 134-9 (1954).—The grains of rye, which had originated in the ears of wheat in the previous generation, have an intermediate chem. compn. between rye and wheat, the examn. being made on the basis of viscosity of aq. ext., glossiness of the grain, quality of flour and bread, and amylolytic activity. G. M. Kasolapoff

KARAPETYAN, V. K.

U S S R .

✓ The surface films of the gliadins of rye with wheat heredity. V. L. Kretovich, G. A. Deborin, A. A. Baudel, L. B. Gorbacheva, and V. K. Karapetyan (A. N. Bakh Inst. Biochem., Acad. Sci. U.S.S.R., and Inst. Genetics, Acad. Sci. U.S.S.R.). *Biokhim. Zerna, Akad. Nauk S.S.S.R., Sbornik 2*; 140-6(1954).--The surface films of rye grains found in wheat ears were studied. The mol. wt. of gliadin from wheat or rye is 30,000. The mol. wt. of the specimens taken from the "altered" grains is but 15,000; this material also shows significantly greater limiting area in formation of a monolayer, than is the case for the normal wheat or rye. Thus, the formation of these grains is accompanied by a severe alteration of the protein structure.

G. M. Kosolapoff

KARAPETYAN, V.K.

New data on plant species formation. Trudy Inst.gen. no.22:
121-136 '55. (MLRA 9:4)
(Wheat)

KARAPETYAN, V.K.

Winter and frost hardiness of soft wheats produced from hard spring
wheat and hard semiwinter wheat. Trudy Inst.gen.no.23:181-189 '56.
(MIRA 10:1)

(Wheat) (Plants--Frost resistance)

KARAPETYAN, V.K.

The problem of species and species formation in the light of studies
on interspecific hybridization of plants. Trudy Inst. gen. no.24:35-75
'58. (MIRA 11:9)

(Hybridization, Vegetable) (Species)

KARAPETYAN, V.K.

Biological effect of nuclear radiation on plants. Agrobiologia
no.1:82-85 Ja-F 60. (MIRA 13:5)

1. Institut genetiki Akademii nauk SSSR.
(Plants, Effect of radioactivity on)

KARAPETYAN, V.K.

Changing winter wheat varieties with poor wintering characteristics
into winter hardy varieties. Agrobiologiya no.6:878-885 N-D '60.
(MIRA 13:12)

1. Institut genetiki Akademii nauk SSSR.
(Wheat)

KARAPETYAN, V.K.

Converting hard frost-sensitive winter wheat and spring wheat
into frost-resistant winter wheat. Trudy Inst. gen. no. 27:54-
67 '60. (MIRA 13:12)
(Wheat) (Plants--Frost resistance)

KARAPETIAN, V.K.

Effect of gamma rays on heredity in wheat. Trudy Inst. gen.
no. 27:311-314 '60. (MIRA 13:12)
(Plants, Effect of gamma rays on)
(Wheat) (Heredity)

KARAPETIAN, V.K.

Effect of gamma rays on soft wheat and rye. Trudy Inst. gen.
no. 27:315-318 '60. (MIRA 13:12)

(Plants, Effect of gamma rays on) (Wheat)
(Rye)

KARAPETYAN, V.K.

Winterhardiness and developmental characteristics of interspecific
hybrid plants. Trudy Inst. gen. no.28:91-95 '61. (MIRA 14:11)
(WHEAT BREEDING) (PLANTS—FROST RESISTANCE)

KARAPETYAN, V.K.

Effect of ionizing radiation on heredity and vitality in some wheat
and rye varieties. Trudy Inst. gen. no.28:141-145 '61.
(MIRA 14:11)

(WHEAT) (RYE)
(PLANTS, EFFECT OF GAMMA RAYS ON)

KARAPETYAN , V.K.

"The effect of ionizing radiation on the heredity and life of some varieties of wheat and rye."

Report submitted to the 2nd Intl. Colloq. on Insect Pathology and
Microbiological Control, Paris, France 16-24 Oct 1962

S/670/62/000/029/004/006
D291/D307

AUTHOR: Karapetyan, V.K.

TITLE: The biological effect of ionizing radiation on variability in spring wheat during its conversion to the winter habit

SOURCE: Akademiya nauk SSSR. Institut genetiki. Trudy.
no. 29, 1962, 194-206

TEXT: In 1957 work was initiated at the Institut genetiki AN SSSR (Institute of Genetics of the AS USSR) on the use of ionizing radiation for shattering the hereditary basis of plants, with subsequent cultivation under different environments. The present investigation was intended to show that when spring wheat is treated in this manner, forms differing in their developmental habit arise. Air-dry seeds of the Armenian spring variety Erinaceum (Triticum compactum var, erinaceum) were treated with Co⁶⁰ gamma rays at dosages ranging from 8 to 16 kr. The 10-16 kr doses were excessive, but a number of plants derived from 8 kr-treated seed survived the

Card 1/3

The biological effect ...

S/670/62/000/029/004/006
D291/D307

winter when sown in autumn and were in fact shown to be winter forms genetically. After two generations of autumn sowing, different families of the progeny of one of the surviving plants showed survival percentages of 37.8 to 82.4%. They were incapable of caring when sown in spring. Subsequently other winter forms were obtained in a similar manner, having greater overwintering capacity than the standard winter wheat variety Ukrainka. The winter selections produced grain yields which were, on average, 10-12% higher than in the original spring form. They were often early maturing, and had a long vernalization stage, not less than 50-55 days. 60% of the winter forms resembled the original variety morphologically. Others were classified as *T. compactum* var. *rubriceps* and *T. vulgare* vars. *ferrugineum* and *erythrospermum*. Some of the winter forms were compact, with strong, lodging-resistant straw; grains were often large, vitreous, uniform and high in protein. These compact forms were genetically stable, retaining their properties in successive generations. It is thought that some of the types which combine earliness, winter hardiness and high productivity will be of practical value. In assessing the results it is pointed out that the treatments applied

Card 2/3

The biological effect ...

S/670/62/000/029/004/006
D291/D307

offer a very rapid way of converting spring into winter wheat, the effects are caused by environmental influences on plants with shattered inheritance. There are 6 figures and 4 tables.

Card 3/3

KARAPETYAN, V.K.

Variability of plants grown from embryonally young seeds.
Trudy Inst. gen. no.29:435-441 '62. (MIRA 16:7)

(Botany—Variation)

KARAPETIAN, V.K. (Karapetyan, V.K.)

Ionizing radiation and its action on the heredity and vitality
of some varieties of wheat and rye. Analele biol 16 no.1:12-18
Ja-F'62

KARAPETYAN, V. K.

"Genetical Analysis of ye-Wheat and Rye-Wheat Hybrids."

Report presented at the 2nd International Wheat Genetics Symposium,
19-24 Aug 63, Lund, Sweden.

KARAPETYAN, V.K.

Effect of ionizing radiation on the variability of spring wheat
during its transformation into winter wheat. Agrobiologia no.1:
107-110 Ja-F '63. (MIRA 16:5)

1. Institut genetiki AN SSSR.
(Botany--Variation) (Wheat) (Plants, Effect of radiation on)

KARAPETYAN, V.K.

Controlled hereditary change of dual-purpose wheat into winter-hardy forms of winter wheat. Trudy Inst. gen. no.30:109-118 '63.(MIRA 17:1)

KARAPETYAN, V.K.

Variability of plants grown from embryonially young seeds.
Zhur. ob. biol. 24 no.5:360-365 S-O '63. (MIRA 17:1)

1. Institut genetiki AN SSSR, Moskva.

KARAPETYAN, V.K.

Controlled heritable transformation of the dual-purpose wheat
into frost-resistant winter forms. Izv. AN SSSR. Ser. biol. no.3:
451-459 My-Je '64. (MIRA 17:5)

1. Institute of Genetics, Academy of Sciences of the U.S.S.R.
Moscow.

KARAPETIAN, V.K.; GYULANYAN, A. Ye.

Biological effect of ionizing radiation on the variability of
spring wheat during its conversion into winter wheat. Izv. AN
Arm. SSR. Biol. nauki 15 no.12:33-43 1962 (MIRA 17:8)

1. Institut genetiki AN SSSR i Leninakanakaya gosudarstvennaya
selektsionnaya stantsiya.

KARAPETYAN, V.K.

Characteristics of the development of wheat-rye and rye-wheat
hybrids. Trudy Inst. gen. no.31:119-125 '64. (MIRA 17:9)

KOLESNIKOV, N.A.; KUBYSHV, N.N.; FEDORENKO, V.G.; KARAPETYAN, V.K.;
UNZHAKOV, M.S.

Intensification of the shaft furnace lead smelting process by
augmenting the oxygen concentration. Tsvet. met. 27 no.12:
33-38 D '64 (MIRA 18:2)

OKUNEV, A.I.; CHUMAREV, V.M.; DONCHENKO, P.A.; KARAPETYAN, V.K.

Accelerating the fuming of slags with the use of oxygen-enriched
air. TSvet. met. 36 no.5:34-41 My '63. (MIRA 16:10)

AKHMETOV, K.T.; DONCHENKO, P.A.; KUBYSHEV, N.N.; VOLKOV, I.P.; KARAPETYAN, V.K.;
YELYAKOV, I.I.; CHIKRIZOV, M.V.; KHOBDABERGENOV, R.Zh.

Modernizing the industrial equipment of lead production and the
growth of labor productivity. TSvet. met. 36 no.7:11-19 J1
'63. (MIRA 16:8)

(Lead industry--Equipment and supplies)

ALEKSEYEVSKIY, V.V.; KARAPETYAN, V.M., inzh.; GRIGORYAN, Ye.B., inzh.

New series of distribution transformers with 160 - 630 kv.-a power rating. Vest. elektroprom. 34 no.4:25-26 Ap '63. (MIRA 16:10)

1. Chlen-korrespondent AN Armyanskoy SSR (for Alekseyevskiy).

KARAPETTYAN, V.M.; AYKAZUNI, G.A.

The TSM-180/10 oil-filled power transformer. Biul.tekh.-ekon.
inform. no.11:40-41 ' 58. (MIRA 11:12)
(Electric transformers)

KARAPETIAN, V. M.

37608. sluchay vistseral'nogo leyshmanioza j vzroslogo v g. jerevane. trudy
in-ta malyarii i med. papazitologii (m-vo zdravookheaneniya arm.ssr)
vyp. 4, 1949, s. 152-56.

SO: Letopis' Zhurnal' nykh Statey, Vol. 37, 1949

SOV/110-59-9-2/22

AUTHORS: Oganyan, R.A., Gukasyan, M.G. and Karapetyan, V.M.
(all Engineers)

TITLE: A New Series of 6 and 10 kV Power Transformers of the
First and Second Frame Sizes

PERIODICAL: Vestnik elektropromyshlennosti, 1959, Nr 9, pp 5-8 (USSR)

ABSTRACT: A new series of general-purpose transformers designated type TSM has been developed. Altogether there are ten different ratings, each being greater than the previous one by a factor of 1.73. The ratings in the first frame size are 20, 35, 60 and 100 kVA and in the second frame size 180, 320 and 560 kVA. In addition, subsidiary ratings of 135, 240 and 400 kVA are made in the second frame size. The high-voltage windings will be for 6 or 10 kV, though 6.3 kV will be supplied for existing installations. Off-load tapping switches of $\pm 5\%$ are provided on the high-voltage side. The ratio of copper to iron loss is 3.5 - 4.0. To reduce deterioration of transformer oil and insulation the maximum temperature rise at the top of the oil has been reduced from 60°C to 55°C, and that of the windings reduced from 70°C in the old standard to 65°C. The insulation, and the clearances in the main insulation, are the same for both 6 and 10 kV

Card 1/3

SOV/110-59-9-2/22

A New Series of 6 and 10 kV Power Transformers of the First and Second Frame Sizes

transformers. Further reduction in these clearances is undesirable because it would impair cooling. The cores will be made of annealed cold-rolled steel grade E330, 0.35 mm thick. The sheets will be insulated with varnish. A number of changes have been made, mainly with the object of reducing weight and size or to facilitate manufacture. A transformer type TSM 100/6-10 is illustrated in Fig 1, alongside an old TM series transformer of the same rating. A transformer type TSM 180/6-10 is illustrated in Fig 2. The three-limb core-type construction is used; the core clamps are made of angle iron and the clamping arrangements are described. An illustration of a typical core and coil assembly for the new series of transformers is given in Fig 5. The core and coils no longer have any attachment to the lid and are fully supported by the tank, so that there is less risk of damage in transport. The high-voltage leads are made of flexible wire and the low-voltage leads of copper strip. The new transformer tanks for 20 and 35 kVA are made of 2 mm sheet steel stiffened by cooling ribs. From 60 kVA and upwards

Card 2/3

SOV/110-59-9-2/22

A New Series of 6 and 10 kV Power Transformers of the First and Second Frame Sizes

tube-cooled tanks are used, the tubes being oval and not round. The tanks themselves are of so-called "oval" section. Even the smallest of the 6 kV transformers have conservators. The other fittings are described. The total losses of the new transformers are 15-25% lower than those of transformers conforming to the existing standard GOST 401-41, also of transformers made by the East German firms TR0 and TUR. Fig 3 compares the weights of oil in the old and new series transformers and in the German transformers. If all Soviet transformers in these ratings were made in the new type the total annual economy of copper would be 120 tons, of steel 900 tons, of transformer oil 4000 tons and of other materials, including insulation, 3900 tons. Tests on the new transformers have confirmed that the design and construction are well chosen. There are 5 figures.

Card
3/3

KARAPETYAN, V. N.

37607. k voprosu o lechenii vistseral'nogo leyshmanioza solyjsur'minom- v ogl: trudy in-ta malyarii i med. parazitologin. (m-vo zdravookhraneniya arm ssr), vyp. 4, 1949, s. 147-51

SO: Letopis' Zhurnal' nykh Statey, Vol. 37, 1949

GAMBARYAN, P.P.; KARAPETIAN, V.S.; AYRUMYAN, K.A.; KAZARYAN, K.G.;
MEZHLUMYAN, S.K.

Ecology of the Armenian vole (*Promethesomys schaposchnikovi* Sat.).
Zool. zhurn. no. 10:5-16 '57. (MIRA 11:7)
(Adzhar--Ineretan Range--Field mice)

ORZHESHKOVSKIY, V.V.; KARAPET'YAN, V.S.; TIMOFEYEV, N.V.

Eye diseases in infectious nonspecific polyarthritis. Sov.med.
23 no.7:44-46 J1 '59. (MIRA 12:11)

1. Iz Sochinskogo nauchno-issledovatel'skogo instituta revmatizma
(dir. - prof.M.M.Shikhov) Ministerstva zdavookhraneniya RSFSR.
(EYE DISEASES complications)
(ARTHRITIS complications)

KECHEK, G.A.; KARAPETYAN, V.S.

Methods for direct determination of preformed ammonia and glutamine
in the trichloroacetic acid filtrate of blood. Vop. biochim. 1:177-
183 '60. (MIRA 14:12)

1. Department of Biochemistry, Academy of Sciences of Armenian
S.S.R., Erevan. (AMMONIA) (GLUTAMINE)
(BLOOD ANALYSIS AND CHEMISTRY)

MANVELYAN, M.G.; BABAYAN, G.G.; GEVORKYAN, S.V.; ASLANYAN, D.G.;
KARAPETYAN, V.TS.

Study of the system $\text{Na}_2\text{SiO}_3 - \text{Ca}(\text{OH})_2 - \text{H}_2\text{O}$ at 25°C and of the conditions of the adsorption of sodium hydroxide on a calcium metasilicate precipitate. Izv.AN Arm.SSR.Khim.nauki 14
no.4:309-317 '61. (MIRA 14:10)

1. Institut khimii Sovnarkhoza Armyanskoy SSR.
(Calcium silicate) (Sodium hydroxide) (Adsorption)

DERMIN, Yu.M.; KHAYELIAN, E.S.; KARAPETIAN, V.S.; OSIPOVA, E.N.; AKOPYAN,
Dzh.A.

Participation of γ -aminobutyric acid in the metabolism of
glutamic and aspartic acids, alanine and glutamine and in
neutralization of ammonia in the brain tissue. Vop. biokhim.
mol. 1:45-59 '64. (MIRA 18:9)

1. Institut Biokhimii AN ArmSSR.

BABAYAN, G.G.; KARAPETYAN, V.TS.

Physicochemical properties of aqueous solutions of sodium and potassium silicates. Part 1: Electric conductance and viscosity of potassium silicate aqueous solutions. Izv.AN Arm.SSR.Khim.nauki 17 no.1: (MIRA 17:4) 29-37 '64.

1. Institut khimii Gosudarstvennogo komiteta tsvetnykh i chernykh metallov SSSR.

Karape + VA, V.V.

3561: Karapetian, V. V., Determination of the pressure of an interference-affected wave on a vertical structure by means of electro-hydrodynamic analogies (in Russian), *Trudn.-i. in-in. osnovnykh i fundamentalnykh*, no. 25, 62-84, 1933; *Ref. Zh. Mekh.* 1936, Rev. 4463.

A detailed description is given of the known electro-hydrodynamic analogy method and its application to the measurement of wave pressures. Measured results are quoted.

For comparison, a pressure diagram on the Salter-Flound method is given; the pressures calculated by the latter are higher than those determined by the author.

S. S. Volt, USSR.

Courtesy *Referativnyi Zhurnal*

Translation, courtesy Ministry of Supply, England

3
4.18

JE
MT

L 29083-66

ACCESSION NR: AP5019208

UR/0056/65/049/001/0007/0000

AUTHOR: Balats, M. Ya.; Karapetyan, V. V.; Kondrat'yev, L. N.; Obukhov, Yu. V. ³⁴35

TITLE: Intensity of nonradiative transitions in Ta and Pu^{239} mesic atoms ¹⁹B

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 49, no. 1, 1965, 7-9

TOPIC TAGS: mesic atom, nonradiative transition, tantalum, plutonium, Mu meson, x ray spectrum

ABSTRACT: This is a continuation of intensity measurements of nonradiative transitions in a number of heavy elements (ZhETF v. 38, 1715, 1960 and v. 39, 1168, 1960), carried out by means of a scintillation γ -spectrometer. The authors investigated the mesic x-ray spectra and have determined the ratio of the intensities of the $2p-1s$ transitions in Ta and Pu^{239} relative to Pb. Some modification was made in the experimental set-up for the measurements with Pu in order to accommodate the large background in the γ -spectrometer counter from the natural radioactivity of Pu^{239} . Preliminary measurements have shown that when the γ -detector is loaded by the Pu activity the γ -ray spectrum from the $2p-1s$ transitions in Pb is displaced towards the hard region by 3--5%, but this shift causes no noticeable error in the experimental results. The fraction of the nonradiative $2p-1s$ transitions was determined by comparison of the γ -spectra obtained with lead and with the materials

Card 1/2

L 1571-66 EWT(m)/EWP(t)/EWP(b) DIAAP/IJP(c) JD/JG

ACCESSION NR: AP5019208

UR/0056/65/049/001/0007/0009

AUTHOR: Balats, M. Ya.; Karapetyan, V. V.; Kondrat'yev, L. N.; Obukhov, Yu. V. ⁴⁵43

TITLE: Intensity of nonradiative transitions in Ta and Pu²³⁹ mesic atoms ¹³

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 49, no. 1, 1965, 7-9

TOPIC TAGS: mesic atom, nonradiative transition, tantalum, plutonium, Mu meson, x ray spectrum

ABSTRACT: This is a continuation of intensity measurements of nonradiative transitions in a number of heavy elements (ZhETF v. 38, 1715, 1960 and v. 39, 1168, 1960), carried out by means of a scintillation γ -spectrometer. The authors investigated the mesic x-ray spectra and have determined the ratio of the intensities of the 2p--1s transitions in Ta and Pu²³⁹ relative to Pb. Some modification was made in the experimental set-up for the measurements with Pu in order to accomodate the large background in the γ -spectrometer counter from the natural radioactivity of Pu²³⁹. Preliminary measurements have shown that when the γ -detector is loaded by the Pu activity the γ -ray spectrum from the 2p--1s transitions in Pb is displaced towards the hard region by 3--5%, but this shift causes no noticeable error in the experimental results. The fraction of the nonradiative 2p--1s transitions was determined by comparison of the γ -spectra obtained with lead and with the materials

Card 1/2

L 1571-66

ACCESSION NR: AP5019208

studied. The number of radiative transitions was 1 ± 0.08 and 0.59 ± 0.06 for Ta and Pu^{239} , respectively. The corresponding fractions of nonradiative transition were therefore 0 ± 0.08 and 0.41 ± 0.06 . In the case of tantalum, a correction was made for the solid angle. The results are consistent with the theoretical assumptions of D. F. Zaretakiy and V. M. Novikov (ZhETF v. 41, 214, 1961). "The authors thank Prof. B. Pontecorvo for suggesting the experiment and for interest in the work." Orig. art. has: 2 figures and 1 table.

ASSOCIATION: Institut teoreticheskoy i eksperimental'noy fiziki GKIAE (Institute of Theoretical and Experimental Physics, GKIAE)

SUBMITTED: 18Nov64

ENCL: 00

SUB CODE: NP NP

NR REF SOV: 002

OTHER: 000

Card 2/2

KARAPETYAN, V.Ye.; MAKSAKOV, B.I.; FEOFILOV, P.P.

Absorption and luminescence of divalent samarium in alkali halide single
crystals. Opt. i spekt. 14 no.3:441-443 Mr '63. (MIRA 16:4)
(Alkali metal halide crystals--Growth) (Samarium) (Luminescence)


L 34416-66	EWT(1)/EWT(m)/T/EWF(1)/ETI	IJP(c)	JD/JG
ACC NR: AF6015441	SOURCE CODE: UR/0051/66/020/005/0918/0920		
AUTHOR: <u>Bakhshiyeva, G. F.; Karapetyan, V. Ye.; Morozov, A. M.</u>			
ORG: none			
TITLE: <u>Optical characteristics of lanthanum¹ sodium molybdate single crystals</u>			
SOURCE: Optika i spektroskopiya, v. 20, no. 5, 1966, 918-920			
TOPIC TAGS: molybdate, lanthanum compound, sodium compound, refractive index, crystal optic property			
<p>ABSTRACT: Large single crystals of $\text{LaNa}(\text{MoO}_4)_2$ whose C axis was parallel to the axis of growth were grown on a seed by pulling from the melt, and their absorption spectra and refractive indices were measured. The absorption spectrum of an $\text{LaNa}(\text{MoO}_4)_2$ crystal taken with SF-4 and IKS-14 spectrophotometers is shown in the figure. It is noted that the absorption spectra are typical of all crystals having a scheelite structure. Refractive index measurements showed that the light ray is "fractionated" on passing through an $\text{LaNa}(\text{MoO}_4)_2$ prism, apparently because the lattice of this binary molybdate is highly disordered. This factor is also thought to cause the relatively broad luminescence lines of Nd^{3+} in $\text{LaNa}(\text{MoO}_4)_2$ and the broad ESR lines of this compound reported by other authors. Authors express their deep appreciation to A. I. Stozharov and P. P. Feofilov for their steady interest and helpful discussions, and</p>			
UDC: 535.321 + 535.341:548.0			
Card 1/2			

L 34416-66

ACC NR: AP6015441

2

to Ye. M. Sychev and I. A. Shube for assistance in the work. Orig. art. has: 1 figure and 1 table.



SUB CODE: 20/ SUBM DATE: 18Oct65/ ORIG REF: 003/ OTH REF: 006

Card 2/2 BLG

KARAPET'YAN, Ye.A.; KRYSHOVA, N.A., zaveduyushchaya.

Role of the protective sleep inhibition in narcolepsy and other forms of
sleep pathology. Trudy Inst.fiziol. 1:381-393 '52. (MLBA 6:8)

1. Sektor organicheskikh nervnykh rasstroystv. (Sleep)

ANDREYEV, B.V.; KARAPETIAN, Ye.A.; MAYOROV, F.P., zaveduyushchiy; KRYSHOVA, N.A., zaveduyushchaya.

Peculiarities of nocturnal sleep in narcolepsy according to data obtained by the activity recorder. Trudy Innt.fiziol. 1:376-380 '53. (MLRA 6:8)

1. Laboratoriya fiziologii i patologii vysshey nervnoy deyatel'nosti (for Mayorov and Andreyev). 2. Sektor organicheskikh nervnykh rasstroystv (for Kryshova and Karapetyan). (Sleep)

KARAPETYAN, Ye. A.

USSR/Medicine - Nervous disorders

Card 1/1 Pub. 86 -- 37/37

Authors : Karapetyan, Ye. A.

Title : Somnambulism

Periodical : Priroda 43/10, page 127, Oct 1954

Abstract : The characteristics of sleepwalking are described and a theory as to the physical cause of this phenomenon is presented. Sleepwalking is considered as a pathological condition of the nervous system and directions are given for its treatment.

Institution : ... *Inst Physiology in Pavlov*

Submitted : ...